

Biliary cystadenoma- A case report

¹Dr Disha Bhandary, Junior Resident, Department of Surgery, AJ Institute of Medical Science And Research Institute, India

²Dr Meghana D, Junior Resident, Department of Surgery,, AJ Institute of Medical Science And Research Institute, India

Corresponding Author: Dr Disha Bhandary, Junior Resident, Department of Surgery, AJ Institute of Medical Science and Research Institute, India

How to citation this article: Dr Disha Bhandary, Dr Meghana D, “Biliary cystadenoma- A case report”, IJMACR- January - February - 2021, Vol – 4, Issue -1, P. No. 254 – 259.

Copyright: © 2021, Dr Disha Bhandary, et al. This is an open access journal and article distributed under the terms of the creative commons attribution noncommercial License 4.0. Which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Type of Publication: Case Report

Conflicts of Interest: Nil

Introduction

Biliary cystadenoma occurs in middle aged women characterized by multilocular cysts with internal septae and mural nodules. Unilocular biliary cystadenomas are rare and are difficult to differentiate from cystadenocarcinoma before surgery. Historically these cystic tumors have been treated by a variety of techniques including aspiration

Fenestration , internal drainage and resection. They cannot be safely differentiated from cystadenocarcinoma before operation and always be considered for resection. Less than 200 cases being reported all around the world. Biliary cystadenomas are slow growing rare benign lesions , are easily respectable with a reported recurrence rate of 90% when the resection is incomplete.

Case report

This report here a 30 year old female who presented in general surgery out patient department of the hospital with pain abdomen since 1 week and abdominal distention since one and half years.

On general examination she was moderately built and nourished with history of weight loss since the past one year. There was a mass in a right hypochondrium extending into epigastrium. The other systemic examinations were normal.

USG examination revealed large multiloculated cystic lesion with thick septations and low level internal echoes measuring 16*13*18 cms involving the right and upper lobes and diagnosis of biliary cystadenoma was made.

Computerized tomography (ct) scan of abdomen revealed enhancement within thin walled

Multi-septated appearance of the liver. Minimum intra-hepatic and extra-hepatic biliary radical dilatations. Biliary radicles are dilated.

The patient was investigated for her liver function and total blood count .they weren't within normal limits. He hemoglobin level was initially 8gm/dl following which patient was transfused 1 pint PRBC, the blood count was repeated and it came up to 10mg/dl. Liver function test was on the lower side, following surgery it was normal.

The patient underwent solitary benign hepatic cystectomy and it was sent for histopathological examination.


Histopathological findings revealed hepatobiliary cystadenoma with mesenchymal stroma. No epithelial proliferation/dysplasia/malignancy seen. Gallbladder histologically unremarkable.

The postoperative course of the patient was uneventful and she has been discharged.

Intraoperative



Histopathology



Discussion

Biliary cystadenoma of the liver are rare benign lesions of the liver that account for less than 5% of solitary cysts of the liver and occur commonly in middle aged women (40-50 years) with preponderance of 4:1 among males, our

patient was relatively younger, she was 30 years old. These tumors present commonly as multiloculated cysts as seen in our patient while unilocular biliary cystadenomas are rare, biliary cystadenomas need to be differentiated from other cystic lesions of the liver like hydatid cysts. These neoplasms are reported in middle aged women as true proliferative epithelial tumors that are large multiloculated with internal septations and surrounded by dense cellular fibrostroma.

References

1. Ishak, K.G., Willis, G.W., Cummins, S.D. and Bullock, A.A., 1977. Biliary cystadenoma and cystadenocarcinoma. Report of 14 cases and review of the literature. *Cancer*, 39(1), pp.322-338.
2. Choi, B.I., Lim, J.H., Han, M.C., Lee, D.H., Kim, S.H., Kim, Y.I. and Kim, C.W., 1989. Biliary cystadenoma and cystadenocarcinoma: CT and sonographic findings. *Radiology*, 171(1), pp.57-61.
3. Korobkin, M., Stephens, D.H., Lee, J.K., Stanley, R.J., Fishman, E.K., Francis, I.R., Alpern, M.B. and Rynties, M., 1989. Biliary cystadenoma and cystadenocarcinoma: CT and sonographic findings. *American Journal of Roentgenology*, 153(3), pp.507-511.
4. Forrest, M.E., Cho, K.J., Shields, J.J., Wicks, J.D., Silver, T.M. and McCormick, T.L., 1980. Biliary cystadenomas: sonographic-angiographic-pathologic correlations. *American Journal of Roentgenology*, 135(4), pp.723-727.
5. Thomas, K.T., Welch, D., Trueblood, A., Sulur, P., Wise, P., Gorden, D.L., Chari, R.S., Wright Jr, J.K., Washington, K. and Pinson, C.W., 2005. Effective treatment of biliary cystadenoma. *Annals of surgery*, 241(5), p.769.