

A descriptive analytical research on cervical malignancy, diagnostic colposcopy, cancer pharmacotherapeutics, immunotherapeutics and co-therapeutic modalities

¹Dr. Moumita Hazra, Associate Professor, Head of Department in Charge, Department of Pharmacology, Mamata Medical College and Hospitals, Telangana, India.

²Dr. Samaresh Kumar Hazra, Consultant Obstetrician and Gynaecologist, Medical Proprietor, Hazra Nursing Home, Hazra Polyclinic and Diagnostic Centre, West Bengal, India.

Corresponding Author: Dr. Moumita Hazra, Associate Professor, Head of Department in Charge, Department of Pharmacology, Mamata Medical College and Hospitals, Telangana, India.

How to citation this article: Dr. Moumita Hazra, Dr. Samaresh Kumar Hazra, “A descriptive analytical research on cervical malignancy, diagnostic colposcopy, cancer pharmacotherapeutics, immunotherapeutics and co-therapeutic modalities”, IJMACR- March - April - 2022, Vol – 5, Issue - 2, P. No. 27 – 36.

Copyright: © 2022, Dr. Moumita Hazra, et al. This is an open access journal and article distributed under the terms of the creative commons attribution noncommercial License 4.0. Which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Type of Publication: Original Research Article

Conflicts of Interest: Nil

Abstract

HPV-associated cancers are typical epithelial malignancies that are difficult to treat when metastatic. Chemotherapy generally consists of combinations of cytotoxic agents, often administered in conjunction with a biological agent. These regimens have limited clinical activity and substantial toxicity, and better treatments are needed. Immunotherapy works through different mechanisms than chemotherapy and has been a breakthrough for the treatment of certain malignancies. Targeting HPV oncoproteins with antigen-specific immunotherapy using therapeutic vaccines are under clinical trials for cervical cancer and metastatic disease treatment. Immunotherapy with PD-1–targeted agents has shown clinical activity in genitourinary and oropharyngeal cancers.

This descriptive analytical research study explores cervical malignancy, and the different aspects of

diagnostic colposcopy, cancer pharmacotherapeutics, onco-immuno-therapeutics and co-therapeutic modalities for a comprehensive cervical cancer treatment.

Keywords: Colposcopy, Cancer Pharmacotherapeutics, Immunotherapeutics, Bevacizumab, Ipilimumab, Cemiplimab, Pembrolizumab, Preventive HPV vaccines, GX-188E therapeutic DNA vaccine, Axal Imogene filolisbac, Adoptive T-cell therapy.

Introduction

According to GLOBOCAN 2018, cervical cancer is the fourth most commonly diagnosed tumor and the fourth cause of cancer death in females, worldwide, but ranks second in both incidence and mortality in lower income countries. HPV-associated cancers are typical epithelial malignancies that are difficult to treat when metastatic. Chemotherapy generally consists of combinations of cytotoxic agents, often administered in conjunction with a biological agent. These regimens have limited clinical

activity and substantial toxicity, and better treatments are needed. Immunotherapy works through different mechanisms than chemotherapy and has been a breakthrough for the treatment of certain malignancies. Targeting HPV oncoproteins with antigen-specific immunotherapy using therapeutic vaccines are under clinical trials for cervical cancer and metastatic disease treatment. Immunotherapy with PD-1-targeted agents has shown clinical activity in genitourinary and oropharyngeal cancers.

Objective

The objective of this descriptive analytical research study was to explore cervical malignancy, and the different aspects of diagnostic colposcopy, cancer pharmacotherapeutics, onco-immuno-therapeutics and co-therapeutic modalities for a comprehensive cervical cancer treatment.

Methodologies, results and Discussion

A Descriptive Analytical Research Study

Oncological diagnostic applications of Colposcopy:

Colposcopy is a diagnostic procedure performed to evaluate women with an abnormal Papanicolaou (Pap) test, women with visual inspection with acetic acid, women positive for high-risk human papillomavirus (HPV) DNA, or with a suspicious appearing cervix even if the PAP test is normal. It is also performed as a post-treatment follow-up of intraepithelial and invasive carcinoma. Colposcopy is a procedure in which a lighted, magnifying instrument called a colposcope is used to examine the cervix, vagina, and vulva. It is a diagnostic procedure performed to evaluate abnormal cytology results from a screening Pap test. It is well known that colposcopy has significant variability and poor reliability between colposcopists. The ASCCP (American Society for Colposcopy and Cervical

Pathology) published colposcopy standards in 2017 to address these concerns related to colposcopy. Colposcopy is a procedure in which a lighted, magnifying instrument called a colposcope is used to examine the cervix, vagina, and vulva. It is a diagnostic procedure performed to evaluate abnormal cytology results from a screening Pap test.

The indications for colposcopy include evaluation of women with an abnormal Pap test to localize the lesion, to map out the extent of the lesion, to select the biopsy site or sites; women positive for high-risk HPV DNA; visual inspection with acetic acid positive women, suspicious appearing cervix, postcoital or postmenopausal bleeding, even if the Pap smear is normal; unexplained abnormal lower genital tract bleeding; persistent inflammatory or unsatisfactory cervical cytology despite appropriate treatment, especially with high-risk factors for carcinoma cervix; persistent abnormal vaginal discharge or pruritus vulvae; for the identification and management of subclinical papillomavirus infection; patients with history of *in utero* diethylstilbestrol (DES) exposure; for the conservative management of intraepithelial neoplasia; identification and management of vaginal extension of cervical neoplasia; and for post-treatment follow-up after treatment of intraepithelial and invasive carcinoma and post-irradiation follow-up.

The ASCCP (American Society for Colposcopy and Cervical Pathology) has published standardization guidelines for the performance of colposcopy. The ASCCP makes recommendations for extensive and minimum requirements for a colposcopy. The colposcopist should examine the vulva, vagina, and cervix grossly in the natural state and also after the application of 5% acetic acid. The entire cervix and

squamocolumnar junction must be visualized for adequacy. Both white light and a red-free (blue or green) filter should be applied to the visual field in order to identify any lesion.

Directed biopsies of lesions should be taken of each abnormal finding. Documentation in a minimum of text format should comment on the visibility extent, size, location, and description of each lesion (color, contour, border, vascular changes), presence or absence of acetowhite Ning, complete or incomplete visibility of the squamocolumnar junction, documentation of biopsies and locations, if an endocervical curettage was performed, and finally the impression of the colposcopy (benign-normal, low grade, high grade or cancer). Application of Monsel's solution or silver nitrate should be applied after the colposcopy is completed, and all biopsies are taken. Grade of cytological abnormality, colposcopic adequacy, visibility, and type of squamocolumnar junction should be documented. The location of lesion, size, and extent of lesion, endocervical, or vaginal extension of the lesion should also be clearly documented. Abnormal colposcopic findings should be described location wise in detail, and colposcopic impression should be made in terms of low grade or high-grade lesion along with Reid's or Swede score. Histopathologic diagnosis should never be made on colposcopy only. According to the 2011 IFCPC Nomenclature, the Swede score is used to score the colposcopic findings and to have uniformity in the reporting system. The total score is 10. Colposcopy is a diagnostic procedure done due to an abnormal cervical screening test or a visible lesion seen on the cervix during an examination. This diagnostic procedure assists with the formulation of a management plan based on the results of the biopsied pathology or lack of results. In

general, all results can either be observed or treated and are based on evidence-based guidelines. Low-grade lesions can be followed up and managed according to ASCCP guideline algorithms. High-grade lesions are treated depending on the patient's age and fertility status. A patient that is pregnant will have their treatment deferred until after delivery unless there is a specific concern for an invasive lesion. Invasive lesions should be referred to an obstetrician and gynaecologist and a gynecological oncologist for treatment options.

The general assessment is performed to determine whether there is adequacy or inadequacy for the reason (e.g., cervix obscured by inflammation, bleeding, scar); to visualize the squamocolumnar junction and to determine whether the visibility is completely visible, partially visible, or not visible; and also to visualize the transformation zone types 1, 2 and 3.

The colposcopy is performed to distinguish between normal and abnormal colposcopic cervical findings, which includes the visualization of whether the original squamous epithelium is mature, columnar epithelium is atrophic or not, whether there is ectopy, metaplastic squamous epithelium, Nabothian cysts or crypt gland openings.

For visualizing abnormal colposcopic findings, the general characteristic features are looked into. The location of the lesion is observed, whether it is inside or outside the T-zone. The location of the lesion is also observed according to the clock position. The size of the lesion is examined, along with the number of cervical quadrants the lesion covers and size of the lesion in the percentage of the cervix. The grading of the lesion is done as follows : in grade 1, minor lesions, there would be thin aceto-white epithelium, irregular, geographic border, fine mosaic and fine punctation; in grade 2,

major lesions, there would be dense aceto-white epithelium, rapid appearance of aceto-whitening, cuffed crypt (gland) openings, coarse mosaic, coarse punctuation, sharp border, inner border sign and ridge sign; in non-specific lesions, there would be leukoplakia (keratosis, hyperkeratosis), and erosion, Lugol’s staining (Schiller’s test) is done to observe whether the tissues are stained or non-stained. If the lesion is suspicious for invasion, there might be the presence of atypical vessels, or associated signs, like, fragile vessels, irregular surface, exophytic lesion, necrosis, ulceration (necrotic), tumor or gross neoplasm. Also, miscellaneous findings might be seen in congenital transformation zone, condyloma, polyp (ectocervical or endocervical), inflammation, stenosis, congenital anomaly, post-treatment consequence, and endometriosis.

The scoring is done as follows:

Score	0	1	2
Aceto uptake	Zero or transparent	Shady, Milky (not transparent; not opaque)	Distinct, opaque white
Margins or surface	Diffuse	Sharp but irregular, jagged, ‘geographical’ satellites	Sharp and even, the difference in surface level, including ‘cuffing’
Vessels	Fine, regular	Absent	Coarse or atypical
Lesion size	<5mm	<5mm 5-15mm or 2	>15mm or 3-4 quadrants/

		quadrants	endocervical undefined
Iodine staining	Brown	Faintly or patchy yellow	Distinct yellow

Overall Swede score

Overall Swede score	Colposcopic prediction of probable histology
0 – 4	Low grade/normal CIN 1
5 – 6	High grade/non-invasive cancer CIN 2+
7 – 10	High grade/suspected invasive cancer CIN 2+

Human papillomavirus associated malignancies and therapeutics

Human papillomavirus (HPV)-associated cancers are common epithelial malignancies, that account for approximately 5% of all cancers worldwide. They occur at varied genitourinary and oropharyngeal anatomic sites. Advanced-stage HPV-associated cancers are difficult to treat. Advanced-inoperable cervical cancer is a challenging entity due to increased percentage of locoregional and distant recurrences. Recurrent cervical cancer not amenable to radical treatment, as well as metastatic disease, are difficult to cure, with a bad prognosis.

Minimally invasive robotic surgery has become an effective surgical technique for the treatment of gynaecologic malignancies, like cervical cancer. Minimally invasive surgery (MIS) is the standard approach to performance of several gynaecologic procedures including hysterectomy, gynaecologic cancer staging procedures, myomectomy, pelvic organs prolapse repair and select adnexal procedures. Robotic-

assisted surgery, a computer-based MIS approach has been adopted widely in the United States and several other countries.

As for the pharmacotherapeutic modalities of cervical cancer treatment, the approved immune-checkpoint inhibitors, the “first generation,” include monoclonal antibodies directed against PD-1 (pembrolizumab, nivolumab, cemiplimab); against PD-L1 (atezolizumab, avelumab, and durvalumab); and against protein CTLA-4 (ipilimumab). Combination chemotherapy and bevacizumab offers some clinical benefit, and anti-programmed death 1 receptor (PD-1) therapy has shown clinical activity, but these malignancies generally are incurable and better treatments are needed. Sequential ipilimumab after chemoradiotherapy is given in curative-intent treatment of patients with node-positive cervical cancer. Anti-tumour activity of cemiplimab as monotherapy or in combination with hypo fractionated radiation therapy, is given in patients with recurrent or metastatic cervical cancer. Pembrolizumab and GX-188E therapeutic DNA vaccine are also emerging cervical cancer treatment options in patients with HPV-16-positive or HPV-18-positive advanced cervical cancer. Pembrolizumab, an IgG4-kappa humanized monoclonal antibody, against the programmed cell death protein 1 (PD-1) receptor, has been approved for the treatment of recurrent or metastatic cervical cancer. Although immune-checkpoint blockade therapy is rapidly altering the treatment landscape in solid tumours, the efficacy of immune-checkpoint blockade therapy with antibodies directed against CTLA-4, PD-1, and PDL-1 in advanced gynaecologic cancers has been limited. The exception has been the PD-1 inhibitor pembrolizumab in microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) advanced

endometrial cancers, highlighted by the recent conditional approval of pembrolizumab in recurrent or metastatic PDL-1 positive cervical cancers and the accelerated approval of pembrolizumab and Lenvatinib in microsatellite stable (MSS) or mismatch-repair proficient (pMMR) advanced endometrial cancer. GX-188E vaccination has been shown to induce human papillomavirus (HPV) E6-specific and E7-specific T-cell responses and cervical lesion regression in patients with cervical precancer. Preventive HPV vaccines and adoptive T-cell therapy, the systemic infusion of therapeutic T cells, are potential emerging cancer treatment modalities. Cellular therapy has shown to mediate the regression of HPV-associated cervical cancer, oropharyngeal cancer, and anal cancer, including durable, complete regression of cervical cancer.

The currently recommended standard of care treatment of cervical malignancy, according to the respective FIGO staging is that, for FIGO Stages IA1 and IA2, type II radical hysterectomy and pelvic lymph node dissection is suggested; for stages IB1 and IIA1, type III radical hysterectomy and pelvic lymph node dissection are suggested; for stages IB2 and IIA2, pelvic external beam radiation therapy, brachytherapy, and cisplatin based concurrent chemotherapy are suggested; for stages IIB and IVA, pelvic external beam radiation therapy, with brachytherapy and cisplatin-based concurrent chemotherapy, and with or without external beam radiation therapy to para-aortic nodes are suggested; and for stages IVB or recurrent disease not amenable to local therapy, paclitaxel, cisplatin and bevacizumab, or, paclitaxel and cisplatin, or, paclitaxel, topotecan and bevacizumab, or, paclitaxel and topotecan, or, paclitaxel and carboplatin, are suggested.

Targeted therapy in cervical cancer

Antiangiogenic therapy targeting the vascular endothelial growth factor (VEGF) and other pathways has improved outcomes in multiple solid tumours. Poor prognosis and early recurrence in cervical cancer has been associated with VEGF expression. Bevacizumab is a recombinant humanised monoclonal immunoglobulin (Ig)-G1 antibody directed against VEGF-A. By inactivating VEGF-A, it blocks signal transduction through VEGFR-1-associated and VEGFR-2-associated pathways. The other targeted therapies, under various phases of clinical trials, include tyrosine kinase inhibitors sunitinib, which inhibits VEGFR, PDGFR, c-KIT, and FLT-3, and pazopanib, which inhibits VEGFR, PDGFR, and c-KIT. Brivanib, an inhibitor of VEGFR and FGFR, is also another targeted therapy modality of advanced cervical cancer, which is under clinical trial. Other VEGF/ VEGFR targeting drugs, like, nintedanib and cediranib are also investigational drugs. Cervical cancer expresses moderate to high levels of epidermal growth factor receptor (EGFR) protein. Several studies with EGFR-targeted therapies, gefitinib, erlotinib and cetuximab, are undergoing different phases of clinical trials. Different clinical trials were also conducted with lapatinib, a HER2 inhibitor, and pazopanib (combination of HER2 inhibitor with VEGFR inhibitor). A recent study of molecular profiling of cervical cancer samples and testing in patient-derived xenograft (PDX) models has demonstrated that co-administration of trastuzumab and lapatinib, the BCAR4, breast cancer anti-estrogen resistance 4 amplification or HER2-overexpressed drugs in PDX significantly inhibited tumour growth compared with the control. A mTOR targeting drug, like temsirolimus, and histone deacetylase (HDAC) targeting

drug, like valproic acid, are also under investigational phase.

PD-1 (programmed cell death 1) and PD-L1 expression on cervical cancer infiltrating T cells and dendritic cells, respectively, has been reported to be associated with high-risk HPV positivity and increasing cervical intraepithelial neoplasia grade. PD-1 is expressed by a high fraction of infiltrating CD8 T cells in cervical cancer, suggesting that blocking of PD-1, by the immune checkpoint inhibitors, like pembrolizumab and ipilimumab, might have therapeutic potential and is undergoing clinical trials. Nivolumab, a fully human antibody against PD-1, is also undergoing different clinical trials as a second-line treatment of cervical cancer. PARP inhibitors, like, olaparib and veliparib, are also being investigated for a potential anti-carcinomatous drug target. Poly (ADP-ribose) polymerase (PARP) is a constitutively expressed enzyme that is involved in base excision DNA repair as well as cell replication, transcription, differentiation and gene regulation, and its inhibition has been shown to be synthetic lethal with homologous recombination DNA repair defects. The PARP inhibitor veliparib was studied in combination with cytotoxic therapy in women with recurrent or persistent cervical cancer after receiving pelvic radiation, with or without cisplatin.

The immune checkpoint inhibitor approach is likely to provide higher benefit in earlier lines of treatment and perhaps in combination with other strategies such as chemotherapy and / or radiotherapy.

Immunotherapeutics of cervical carcinoma

The rationale for immunotherapy in cervical carcinoma (CC) is that, given that almost all CCs are human papillomavirus (HPV)-related tumours, CC could represent a paradigmatic example for the benefit

obtained from immunotherapy. The immune system is often stimulated by non-human (viral) antigens, and for this reason it was possible to develop a vaccine as tumour prophylaxis. Several studies have confirmed that a large number of genomic alterations are found in CC patients, for example, in the following genes: KRAS, PIK3CA, TP53 and PTEN. This high mutational burden might be responsive to immunotherapy. HPV-integrated genes are often described in CCs.

In a pharmacogenomic study, it was identified that 384 integrated gene sites could influence T cell activation in the KEGG (Kyoto Encyclopedia of Genes and Genomes, <https://www.genome.jp/kegg/>) database, indicating the possibility of a strong correlation between HPV infection and immune surveillance. There is an interesting correlation between HPV-mediated immune tolerance and tumour development. The ability of HPV to promote a so-called “non-lytic life cycle” inactivates (or partially activates) dendritic cell migration to lymph nodes and consequently inhibits immune activation. At the same time, low expression of E6 and E7 HPV proteins reduces Langerhans cell activity, leading to an immune-tolerant status that can potentially promote CC development. With regard to immune checkpoints, high levels of CTLA4 and PD1/PD-L1 are often detected in CC patients, and PD1/PD-L1 are frequently expressed in dendritic cells in Cervical Intraepithelial Neoplasia (CIN) samples. PD1/PD-L1 expression has been shown to be present in 95% of intraepithelial lesions and around 80% of squamous carcinomas. Several studies on CC have demonstrated high expression levels of immune-suppressive cytokines, such as IL-10, confirming an interesting link between immune checkpoints and CC progression. Recently, it was shown that PD-L1

expression correlates with TILs, predicting response in CC patients treated with neoadjuvant chemotherapy.

One antibody–drug conjugate, tisotumab–vedotin, has been studied in cervical cancer patients.

The immunotherapeutic modalities of cervical cancer treatment include the following

A. Prophylactic

- Preventive HPV vaccines

B. Therapeutic

- GX-188E therapeutic DNA vaccine
- Axalimogene filolisbac ADXS11-001
- Adoptive T-cell therapy
- Cervical cancer therapeutic vaccines aim to eradicate HPV-infected cells by stimulating cytotoxic T cells against the viral/tumour antigens. The HPV E6 and E7 oncoproteins are expressed in HPV-associated cancers and are ideal targets for a therapeutic vaccine. Many live bacterial vectors have been explored in HPV therapeutic vaccines including *Listeria monocytogenes*, *Lactobacillus lactis*, *Lactobacillus plantarum*, *Salmonella enterica* and BCG. *Listeria monocytogenes* has the ability to replicate in the cytosol of antigen-presenting cells and infects monocytes and macrophages, allowing bacterial peptide antigens to be processed and presented via both Major Histocompatibility Complex class I and II pathways, generating potent CD8 and CD4 T cell–mediated immune responses. The sensitivity of *Listeria* to antibiotics allows the vector to be killed in vivo as required. The *Listeria*-based vaccine potency is further enhanced by encoding recombinant proteins composed of HPV E6 and E7 antigens fused to immunostimulatory molecules. Axalimogene filolisbac (ADXS11-001), a live, attenuated *Listeria monocytogenes* bacterial vector secreting HPV-16 E7 fused to listeriolysin O (LLO), is

under investigation for treatment of HPV-associated malignancies including cervical cancer. A phase II study evaluated the safety and efficacy of ADXS11-001, administered with or without cisplatin, in patients with recurrent or refractory cervical cancer following prior chemotherapy and / or radiotherapy.

Therapeutic T cells, is an emerging cancer treatment modality that can induce complete tumour responses in some patients with B-cell malignancies or metastatic melanoma. A study was conducted to test whether adoptive T-Cell therapy could mediate the regression of HPV-associated epithelial cancers. A method was established in the study to generate independent tumour-infiltrating lymphocyte (TIL) cultures from fragments of a resected metastatic tumour deposit. Because HPV-associated cancers constitutively expressed the HPV E6 and E7 oncoproteins, immunologically foreign viral proteins that are attractive targets for immunotherapy and cultures with HPV-oncoprotein reactivity were selected preferentially for administration to patients. A completed clinical trial was presented with long-term follow-up, in which, 18 patients with metastatic cervical cancer and 11 patients with other cancers participated. The trial was a phase II design with two cohorts, cervical cancers and noncervical cancers. Cell infusion was preceded by a lymphocyte-depleting conditioning regimen and followed by systemic high-dose aldesleukin.

Tumour responses occurred in 5 of 18 (28%) patients in the cervical cancer cohort and 2 of 11 (18%) patients in the noncervical cancer cohort. Two of the responses in cervical cancer were complete and the ongoing treatment continued for 67- and 53-months follow-up. Responses in the noncervical cancer cohort were in anal cancer and oropharyngeal cancer. The HPV reactivity of the infused

T cells correlated with clinical response. Peripheral blood repopulation with HPV-reactive T cells also correlated with clinical response. These findings supported the concept that cellular therapy can mediate the regression of epithelial cancers, and they suggested the importance of predictive biomarkers and novel treatment platforms for more effective therapies.

In another similar study on the target antigens in two patients, complete responses were experienced in the clinical trial. Tumour-infiltrating lymphocyte therapy was administered to each patient, which targeted HPV antigens. However, the predominant target antigens were a cancer germline antigen in one patient and mutant neoantigens in another patient.

A number of biological agents modulating different signal transduction pathways are currently in clinical development, like cell cycle inhibitors, histone deacetylases, cyclooxygenase-2 (COX-2), mammalian target of rapamycin (mTOR), heat shock protein (HSP), WEE1, NOTCH signalling and others. With a better understanding of the central role of HPV infection in tumorigenesis of cervical cancer, more studies are evaluating the role of immune-directed therapies in cervical cancer, in adjuvant as well as metastatic settings.

Conclusion

Therefore, this descriptive analytical research on cervical malignancy, diagnostic colposcopy, cancer pharmacotherapeutics, immunotherapeutics and co-therapeutic modalities, well-delineates a detailed diagnostic colposcopy, cancer pharmacotherapeutics, immunotherapeutics and co-therapeutic modalities. This study would certainly remain a landmark on the way towards future innovations in newer drug discovery and development of more appropriate, efficacious, safe, high

quality, easily accessible, inexpensive, easily administered and comprehensively suitable drug, for a much better pharmacotherapy and immunotherapy of cervical malignancy, along with the other therapeutic modalities.

Acknowledgements

My gratitude to the Departments of Pharmacology, Clinical Pharmacology, Molecular Pharmacology, Pharmacogenomics, Rational Pharmacotherapeutics, Evidence-Based Medicine, Medical and Reproductive Endocrinology, Obstetrics and Gynaecology, Clinical Oncology, Clinical Pathology, Pathology, Molecular Diagnostics, Molecular Medicine, Clinical Medicine, Dr. Moumita Hazra's Polyclinic And Diagnostic Centre, Hazra Nursing Home, Hazra Polyclinic And Diagnostic Centre, Institute of Post-Graduate Medical Education and Research and S. S. K. M. Hospital, R. G. Kar Medical College and Hospital, Mamata Medical College and Hospitals, Hi-Tech College of Nursing, and Mahuya Diagnostic Centre and Doctor's Chamber, for the successful completion of this research article publication..

References

1. Wentzensen N, Massad LS, Mayeaux EJ, Khan MJ, Waxman AG, Einstein MH, Conageski C, Schiffman MH, Gold MA, Apgar BS, Chelmow D, Choma KK, Darragh TM, Gage JC, Garcia FAR, Guido RS, Jeronimo JA, Liu A, Mathews CA, Mitchell MM, Moscicki AB, Novetsky AP, Papasozomenos T, Perkins RB, Silver MI, Smith KM, Stier EA, Tedeschi CA, Werner CL, Huh WK. Evidence-Based Consensus Recommendations for Colposcopy Practice for Cervical Cancer Prevention in the United States. *J Low Genit Tract Dis.* 2017; 21(4): 216-222.

2. 2019 ASCCP Risk-Based Management Consensus Guidelines for Abnormal Cervical Cancer Screening Tests and Cancer Precursors: Erratum. *J Low Genit Tract Dis.* 2020; 24(4): 427.

3. Quaas J, Reich O, Küppers V. Explanation and Use of the Rio 2011 Colposcopy Nomenclature of the IFCPC (International Federation for Cervical Pathology and Colposcopy): Comments on the general colposcopic assessment of the uterine cervix: adequate/inadequate; squamocolumnar junction; transformation zone. *Geburtshilfe Frauenheilkd.* 2014; 74(12): 1090-1092.

4. Alan M, Gunyeli I, Gultekin M, Sancı M, Yuce K. Correlation of Swede score colposcopy scoring system and histopathological results in patients with high-risk HPV infection other than HPV16 and 18. *Int J Gynecol Cancer.* 2020; 30(1): 35-40.

5. Khan MJ, Werner CL, Darragh TM, Guido RS, Mathews C, Moscicki AB, Mitchell MM, Schiffman M, Wentzensen N, Massad LS, Mayeaux EJ, Waxman AG, Conageski C, Einstein MH, Huh WK. ASCCP Colposcopy Standards: Role of Colposcopy, Benefits, Potential Harms, and Terminology for Colposcopic Practice. *J Low Genit Tract Dis.* 2017; 21(4): 223-229.

6. Massad LS, Einstein MH, Huh WK, Katki HA, Kinney WK, Schiffman M, Solomon D, Wentzensen N, Lawson HW, 2012 ASCCP Consensus Guidelines Conference. 2012 updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors. *Obstet Gynecol.* 2013; 121(4): 829-846.

7. Mayeaux EJ, Novetsky AP, Chelmow D, Choma K, Garcia F, Liu AH, Papasozomenos T, Einstein MH. Systematic Review of International Colposcopy Quality Improvement Guidelines. *J Low Genit Tract Dis.* 2017; 21(4): 249-257.

8. Stevanovic S, Helman SR, Wunderlich JR, Langan MM, Doran SL, Kwong MLM, Somerville RPT, Klebanoff CA, Kammula US, Sherry RM, Yang JC, Rosenberg SA, Hinrichs CS. A phase II study of tumor-infiltrating lymphocyte therapy for human papillomavirus-associated epithelial cancers. *Clin Cancer Res.* 2019; 25(5): 1486-1493.

9. Mayadev J, Brady WE, Lin YG, Da Silva DM, Lankes HA, Fracasso PM, Ghamande SA, Moore KN, Pham HQ, Wilkinson KJ, Kennedy VA, Aghajanian C, Koh WJ, Monk BJ, Schilder RJ. A phase I study of sequential ipilimumab in the definitive treatment of node positive cervical cancer: GOG 9929. *J Clin Oncol.* 2017; 35(15 suppl): 5526.

10. Youn JW, Hur S-Y, Woo JW, Kim Y-M, Lim MC, Park SY. Pembrolizumab plus GX-188E therapeutic DNA vaccine in patients with HPV-16-positive or HPV-18-positive advanced cervical cancer: interim results of a single-arm, phase 2 trial. *Lancet Oncol.* 2020; 21(12): 1653-1660.

11. Mattson JN, Bender DP. Minimally invasive robotic surgery for gynaecologic cancers: A review. *Clin Obstet Gynecol.* 2020; 63(1): 24-29.

12. Vora C, Gupta S. Targeted therapy in cervical cancer. *ESMO Open.* 2019; 3: e000462.

13. Lontos M, Kyriazoglou A, Dimitriadis I, Dimopoulos M-A, Bamias A. Systemic therapy in cervical cancer: 30 years in review. *Crit Rev Oncol/Hematol.* 2019; 137: 9-17.

14. Schepisi G, Casadei C, Toma I, Poti G, Iaia ML, Farolfi A, Conteduca V, Lolli C, Ravaglia G, Brighi N, Altavilla A, Martinelli G and Giorgi UD. Immunotherapy and its development for gynecological (ovarian, endometrial and cervical) tumors: from

immune checkpoint inhibitors to chimeric antigen receptor (CAR)-T cell therapy. *Cancers.* 2021; 13: 840.

15. Rubinstein MM, Makker V. Optimising immunotherapy for gynecologic cancers. *Curr Opin Obstet Gynecol.* 2020; 32(1): 1-8.